

Beschrieben von

Dr. H. D o h r n.

1. *Coptocheilus Sumatranus*. T. perforata, turrita, solidula, striata, sub epidermide olivacea rubella; sutura simplex, impressa; anfr. 7—8 convexi, ultimus penultimo angustior, antice ascendens, basi rotundatus; apertura subcircularis; perist. subduplicatum, album, internum continuum, margine columellari vix inciso, externum ad anfractum penultimum breviter interruptum, plane expansum, margine columellari recedente, ad incisuram incrassatum ed angulatim expansum.

Long. 19—24, diam. 8—9, ap. diam. 8—9 mm.

Habitat in insula Sumatra, Singalang.

2. *Pupina rufilabris*. T. ovato-acuminata, solidula, glaberrima, rubro-cornea; anfr. 5 convexiusculi, ultimus ventrosus, spiram superans; apertura fere verticalis, subcircularis; paries aperturalis lamina transversali, compressa, intrante munitus; perist. laete rufum, vix expansum, incrassatum, margine dextro retrorsum ascendentem, parum arcuato, basali vix protracto, sinistro canali arcuato, sat longo, breviter ascendentem dissecto, supra canalem lato, ad columellam torulose incrassato.

Long. 10, diam. 6, ap. c. perist. long. $5\frac{1}{4}$, lat. 4 mm.

Habitat cum praecedente.

3. *Pupina turgidula*. T. conico-ovata, glaberrima, fulvo-cornea; sutura marginata; anfr. 5 convexiusculi, ultimus ventrosus, spira brevior; apertura verticalis; paries aperturalis lamina verticali, basi triangulari dein compressa, breviter intrante munitus; perist. expansiusculum, vix incrassatum, concolor; margine dextro pone laminam recte ascendente, dein regulariter arcuato, basali non protracto, ad columellam breviter transverse inciso, area columellari percallosa, dilatata.

Long. 5, diam. 3, ap. c. perist. long. $2\frac{1}{4}$ mm.

Habitat cum praecedente.

4. *Parmella elongata*. T. oblonga, tenuissima, subtiliter striata, superne convexiuscula, nitens, pallide cornea; spira parva, plana; anfr. $2\frac{1}{2}$ subtus aperti, ultimus marginibus dextro et postico membrana tenuissima circumdatus; apert. totam testam occupans.

Diam. maj. 12, min. $7\frac{1}{2}$, alt. 3 mm.

Habitat cum praecedente.

5. *Nanina Sarawakana*. T. perforata, sinistrorsa, acute carinata, depresso turbinata, pellucida, albido-cornea, ad peripheriam rufo-fusco bifasciata; spira convexo-

conoidea, apice obtusula; anfr. 6 subplani, lente accrescentes, primi 2 vix striati, ceteri confertim et subtiliter decussati, ultimus infra carinam distinctius striatus, nitens; apert. diagonalis, oblique securiformis; perist. simplex, rectum, margine columellari circa perforationem protracto.

Diam. maj. 19, min. 17, alt. 8 mm.

Habitat in insula Borneo: Sarawak.

6. *Helix Doriae*. T. semiobtecte umbilicata, turbinatoglobosa, solida, non nitens, obliqua striata, irregulariter malleata, rufo-fusca vel castaneo-fusca, peripheria plerumque nigro vel fusco et albo fasciata; spira convexa apice obtusa; anfr. 5, superi convexiusculi, ultimus ventrosus, peripheria subangulatus, antice descendens; apertura truncato-ovalis, perobliqua, intus margaritacea; perist. undique incrassatum expansum et reflexum, nigrum, margine columellari circa umbilicum dilatato.

Diam. maj. 35—38, min. 29—32, alt. 25—30, ap. lat. 22—26 mm.

Habitat in parte boreali insulae Borneo.

Diagnosen neuer Mollusken.

Von

Dr. A. Westerlund, Ronneby.

Daudebardia haliciensis nov. sp.

Testa umbilicata, umbilico ad aperturam sat forte dilatato, oblongo-ovata, diaphana, hyalino-virescens, distanter tenue striata; anfr. $3\frac{1}{3}$, primi regulariter accrescentes, ultimus forte accrescens, antice dilatatus; sutura marginata; spira fere dimidiata testae efficiens; apertura obovata (extus latior), margine exteriore leviter arcuata, testam non superante, columellari